
Invited Editorial

African Pygmies, What's Behind a Name?

PAUL VERDU^{1*} AND GIOVANNI DESTRO-BISOL^{2,3}

Numerous populations spread out across Central Africa were named Pygmies by Western explorers, since the 19th century, in reference to the mythical Pygmy population from Homer (“A people of short stature at war with migrating birds,” *Iliad*, song 3, v. 1–6). For more than 40 years, Pygmy populations have attracted the interest of biological anthropologists because they are seen as a lively example that the adaptation to the environment shapes the diversity of external physical or physiological characters (e.g., Campbell and Tishkoff 2010; Froment 1993; Perry and Dominy 2009).

Some biological anthropologists and geneticists wrongly perceived Central African Pygmies as a homogeneous group with similar cultural, morphological and biological features, and somewhat opposed to non-Pygmy populations (often designated as “Bantu populations,” though they do not necessarily speak Bantu languages which Pygmy populations often speak). This view may be seen as the consequence of two misleading notions widespread in Western culture. First, the prevalent historical use of the single exogenous term “Pygmy” to designate more than twenty human groups from the Congo Basin suggests that common historical, cultural, and even biological or morphological features are shared by Central African Pygmies. Second, Pygmy populations are often viewed as being exclusively forest hunter-gatherers, isolated from non-Pygmy farmers, and living in the same way as ancestral human populations did before the Neolithic revolution.

Since the 1960s, scientific research has repeatedly challenged such stereotypes. For instance, cultural anthropologists observed that the various Pygmy populations do not share a common myth of origin. Most of them do not know each other and are unaware of a common designation as “Pygmies” from

¹Department of Biology, Stanford University, Stanford, CA 94305, USA.

²Dipartimento di Biologia Ambientale, Università di Roma “La Sapienza.”

³Istituto Italiano di Antropologia, Roma.

*Correspondence to: Paul Verdu, Department of Biology, Stanford University, 371 Serra Mall, Stanford CA 94305, USA. Tel: 1-650-724-5122, Fax: 1-650-724-5114. E-mail: verdu@stanford.edu.

Human Biology, February 2012, v. 84, no. 1, pp. 1–10.

Copyright © 2012 Wayne State University Press, Detroit, Michigan 48201-1309

KEY WORDS: AFRICAN PYGMY, GENETIC DIVERSITY, BIOLOGICAL ANTHROPOLOGY, CULTURAL ANTHROPOLOGY, PYGMY STATURE, LINGUISTICS, EVOLUTION, ADAPTATION, INTERDISCIPLINARY PERSPECTIVES.

outsiders; in fact, the various Aka, Baka, Bongo, Efe (...) Pygmy populations are not a federated group of populations (Bahuchet 1993; Hewlett 1996). Additionally, physical anthropologists have observed a wide diversity of average height across African Pygmy populations and found no height discontinuity between Pygmies and non-Pygmies—some neighboring non-Pygmy populations also are of short stature (Froment 1993; Hiernaux 1974). Finally, highlights are: Pygmies are not exclusively hunter-gatherers—they often practice fishing and agriculture (e.g., Bahuchet et al. 1989; Bahuchet 1992; Hewlett 1996)—and Pygmies are not isolated from neighboring non-Pygmy populations—these groups of populations have close socio-economic relationships including intermarriages (e.g., Bahuchet and Guillaume 1982; Hewlett 1996; Joiris 2003; Kazadi 1981; Turnbull 1965).

The Short Stature of African Pygmies. The issues raised by the common view that Pygmies form a uniform group are best (but not only) exemplified by extensive research focusing on the short stature of these populations. Three “classical” hypotheses were proposed to explain this phenotype: an adaptation to the hot and humid environment of the rainforest (Cavalli-Sforza 1986), an adaptation to the difficult mobility in a densely forested environment (Diamond 1991), or, finally, the outcome of the scarcity of food resources in the forest (Bailey et al. 1989). A fourth hypothesis was recently proposed: The short stature of Pygmy populations could be an adaptation to the earlier onset of reproductive life in a context of high rates of mortality from parasitism and poor nutrition (Migliano et al. 2007).

While being relevant approaches to the understanding of Pygmy height, each one of these hypotheses fails to explain the variability of adult stature observed today within and among the various Central African Pygmy populations when considered separately. Nor can such theories account for the differential stature between specific Pygmy and neighboring non-Pygmy populations that live in comparable climatic, nutritional, or parasitic environments (Becker et al. 2010; Froment 1993; Perry and Dominy 2009).

In fact, understanding the evolution of stature differences across Central African populations requires a tremendous effort in gathering much more anthropometrical, physiological, and demographic data than those currently available and previously used. More specifically, one must gather longitudinal data in the various populations designated as Pygmies instead of taking a single Pygmy group (e.g., the widely studied Aka from Central African Republic) as being representative of all Pygmies across Central Africa (Becker et al. 2010). With richer data taking into account the cultural, environmental and morphological diversity of African Pygmies, anthropologists could better understand the respective roles of the climatic, environmental, and demographic forces that shaped the differential stature among and between the various Pygmy and non-Pygmy populations from Central Africa.

Despite archeological remains attesting to the dense and continuous presence of *Homo sapiens* in the Congo Basin since at least 40,000 years

(Cornelissen 2002), no evidence exists of an ancient *Homo sapiens* population of a particularly short stature having lived in Central Africa. Consequently, we do not know if Pygmy populations evolved toward shorter statures or if their non-Pygmy neighbors evolved toward taller statures. To this end, knowing whether Central African Pygmy populations have an independent or a common origin is an essential starting point, which was somewhat overlooked by some biological anthropologists. If Pygmy populations had independent origins, then their short stature could be the result of convergent evolution while, if they had a recent common origin, their short stature was possibly inherited and the adaptation process (if any) would have occurred in a single ancestral population sometime in the past. If an accurate evolutionary framework and a more representative sampling methodology are to be developed concerning the evolution of Central African populations, biological anthropologists and geneticists should consider the complexity and the ambiguities raised by the binary categorization of populations in Pygmies and non-Pygmies in the multi-ethnic context of this region.

A Glimpse into the Evolutionary History of African Pygmies: A Change of Perspective. In the book *African Pygmies* (1986), Luigi Luca Cavalli-Sforza reviews the results of demographic, medical, genetic, and anthropological researches initiated in 1966. According to this scholar, the Sua (or Mbuti) Pygmies from the Ituri forest would be directly descended from the ancestors of most—if not all—Pygmy groups, whereas the Aka from the Central African Republic would have undergone a high level of genetic admixture with non-Pygmy neighboring populations. According to such results and while being unable to provide formal evidence, Cavalli-Sforza favored the hypothesis of a common Pygmy origin with subsequent diversification, as opposed to the theory of Hiernaux (1974) of independent adaptations to the rain-forest environment.

More than 15 years after the publication of *African Pygmies*, the genetic admixture pattern initially found in the Aka Pygmies was further analyzed by using sex-specific genetic markers (mitochondrial DNA and Y-chromosome polymorphisms), and signatures of a sex-biased genetic admixture between Aka Pygmy groups and non-Pygmies were detected by Destro-Bisol et al. (2004b), in agreement with the ethnographical origins of sex-biased matrimonial regimes. The same research team further hypothesized a common origin of all Pygmy populations and suggested, on the basis of simulation approaches and on phylogenetic inferences, that a first separation between the ancestors of Pygmy and non-Pygmy populations took place between 60,000 and 30,000 years ago and suggested a more recent divergence between the two Pygmy groups of populations now living at the longitudinal extremes of Central Africa with a split occurring prior to 18,000 years ago (Batini et al. 2007; Destro-Bisol et al. 2004a). Such estimates were later confirmed by a phylogenetic approach using

mitochondrial DNA sequences on a wider population set (Quintana-Murci et al. 2008). However, these studies could not formally test the common or independent origin of Central African Pygmies by using a model-based approach since they considered a very limited number of Pygmy population and/or only two non-recombining sex-specific loci. These limitations have since been overcome.

Indeed, starting with 2009, fundamental advances concerning the reconstruction of the history and diversity of Central African populations were reached thanks to the high-resolution population sampling obtained by interdisciplinary teams of biological and cultural anthropologists. In this context, the population-sampling strategy of Verdu et al. (2009) benefited from the numerous cultural criteria suggested by cultural anthropologists (e.g., Bahuchet 1992; Hewlett 1996; Joiris 2003). Verdu et al. (2009) collected more than 600 DNA samples from nine Pygmy populations living in Cameroon and Gabon and from twelve non-Pygmy neighboring populations with whom Pygmies maintain close socio-economic interactions. Their analysis enabled (1) the quantification of the extent of autosomal neutral genetic diversity in Western Central African populations, (2) the investigation of the complex Pygmy/non-Pygmy categorization from a genetic perspective, and (3) the formal test of a common or independent origin of numerous Western Central African Pygmy populations through a model-based approach.

Far from being a genetically homogeneous group of populations, the genetic diversity of Western Central Africans is mainly explained by considerable genetic distances between pairs of Pygmy populations, non-Pygmy populations often being not significantly genetically different from each other (Figure 1). This biological diversity echoed the large cultural diversity observed in the populations sampled and further challenged the common origin conveyed by the use of the blanket term “Pygmy.”

Interestingly, the linguistic diversity of Western Central African Pygmy and non-Pygmy populations did not predict the genetic distances computed among pairs of populations and non-Pygmy neighbors. While speaking closely related languages, they can be genetically very distant from each other (e.g., Baka Pygmies and Bangando non-Pygmies, see Figure 1). Differently, non-Pygmy populations speaking languages belonging to different linguistic families have small, often not significant, genetic distances (see Figure 1).

Furthermore, each Pygmy population had different levels of genetic admixture with its non-Pygmy neighbors (Patin et al. 2009; Tishkoff et al. 2009; Verdu et al. 2009). These heterogeneous levels of asymmetrical gene flow from non-Pygmies into Pygmies were found inversely related to the strength of socio-cultural barriers that impede intermarriages between each specific pair of neighboring communities (Verdu et al. 2009). These results illustrate not only how the diversity of genetic patterns across Pygmy populations is determined by specific socio-cultural factors, but also how the description of the biologic diversity of Central African Pygmies can be biased when only a few groups are analyzed.




Figure 1. Metric multidimensional scaling (MDS) accounting for genetic dissimilarities between central African Pygmy and non-Pygmy populations based on F_{ST} measures (Weir and Cockerham 1984) obtained by analyzing 28 autosomal tetranucleotide microsatellite markers genotyped in 9 Pygmy and 12 non-Pygmy neighboring populations (F_{ST} matrix previously published in Verdu et al. 2009). Non-significant pairwise F_{ST} values based on permutation tests were set to 0 prior to computing the metric MDS (Verdu et al. 2009). To evaluate how accurately the two-dimensional MDS plot represented the pairwise population genetic dissimilarities, we computed the Spearman correlation between the Euclidian distances among pairs of populations on the MDS plot and the corresponding values of pairwise genetic F_{ST} . We obtained a Spearman's rho equal to 0.84 (P -value $< 10^{-5}$). Furthermore, we found that the Euclidian distance calculated between any two populations on the MDS plot provided the corresponding F_{ST} value plus or minus 0.004 on average. Pygmy populations are labeled in black as follows: BEZ = Bezan, central Cameroon; CBK = Baka, central-eastern Cameroon; EBG = Bongo, eastern Gabon; EBK = Baka, southeastern Cameroon; GBK = Baka, northern Gabon; KOL = Kola, western Cameroon; KOY = Koya, northeastern Gabon; SBG = Bongo, southern Gabon; SBK = Baka, southern Cameroon. Non-Pygmy populations are labeled in grey as follows: AKL = Akele, eastern Gabon (Bongomo); BGD = Bangando, southeastern Cameroon; CFG = Fang, southern Cameroon; EWD = Ewondo, central Cameroon; GFG = Fang, northern Gabon; KOT = Kota, central Gabon; NGB = Ngumba, western Cameroon; NZE = Nzebi, southeastern Gabon; NZI = Nzime, central-eastern Cameroon; TEK = Teke, eastern Gabon; TIK = Tikar, central Cameroon; TSG = Tsogho, central Gabon. The linguistic affiliation of samples is mainly based on Greenberg (1966) and Guthrie (1967) and reads as follows: Adamawa-Ubangian family = (▲); Bantoid Non-Bantu family = (+); Bantu family: A. 70 = (▽); A. 80 = (●); B. 20 = (■); B. 30 = (⊕); B. 30-B. 50 = (▽); B. 50 = (⊕); B. 70 = (◆).

Toward the quest of the genetic determination of the differential stature between Pygmies and non-Pygmies, the work of Becker et al. (2011) is very promising. They have shown that Pygmy individuals that are more admixed with non-Pygmy neighbors are also likely to be taller. By analyzing the patterns of genetic admixture across several Western Central African Pygmy populations, they demonstrated, though indirectly, that the differential stature phenotype observed between Pygmies and non-Pygmies had a genetic basis.

Finally, using genome-wide autosomal microsatellites and Approximate Bayesian Computation (ABC) methods (Beaumont et al. 2002; Cornuet et al. 2008), Verdu et al. (2009) formally tested, for the first time, whether numerous Western Central African Pygmy populations shared a common or an independent origin, and estimated the divergence times between populations. Despite the wide genetic differentiation among Western Central African Pygmies, the most probable scenario appeared to be that of a very recent split of Western Central African Pygmies, about 3000 years ago, from an ancestral Pygmy population that diverged from non-Pygmy lineages 50,000 to 90,000 years ago (Verdu et al. 2009). This latter estimate overlaps a previous estimation based on simulation and phylogenetic approaches by Destro-Bisol (2004a; 2004b), Batini et al. (2007), and Quintana-Murci et al. (2008).

A similar model-based methodology was later extended to other Central African Pygmy groups, particularly in the East, by using neutral autosomal sequences (Patin et al. 2009) and mitochondrial DNA (Batini et al. 2011). Results showed that Eastern and Western Central African Pygmy populations, today living several thousand kilometers apart, probably shared a common origin some 20,000 years ago, while the divergence between the ancestral Pygmy and non-Pygmy populations occurred around 70,000 years ago. Both estimates show confidence intervals that are consistent with previous studies based on simulation, phylogenetic, and model-based approaches (Batini et al. 2007; Destro-Bisol et al. 2004a,b; Quintana-Murci et al. 2008; Verdu et al. 2009).

The studies we mentioned so far addressed the diversity of Central African populations in the framework of the complex cultural categorization Pygmy/non-Pygmy and did shed light on the widely unknown evolutionary and adaptation history of Central African populations. We would like to stress here that this research was only made possible by a proficient collaboration between biologists, cultural anthropologists, and linguists. Future multidisciplinary work will allow the careful reconstruction of the history of the admixture between Pygmies and non-Pygmies, as we do not know how the dynamic of intermarriages between Pygmies and non-Pygmies has evolved through history. More specifically, we would like to assess whether genetic admixture patterns have changed over pre-colonial, colonial, and post-colonial times in order to understand how changes in the rates of gene flow may have influenced the biological and genetic adaptation processes of these populations.

Beyond Genetics. We were asked to write this editorial summarizing ongoing genetic research in Central Africa to supplement the article by Serge Bahuchet (Bahuchet 2012, same issue) about the linguistic diversity of Pygmy populations and their non-Pygmy neighbors. Throughout his paper, Serge Bahuchet outlines at least two important messages to biological anthropologists and human biologists, the importance of which goes beyond the specific field of studies on African Pygmies.

First, he notes that the questions asked by linguists are often complementary to those that are the focus of molecular anthropologists. Such questions highlight the importance of the understanding of the social dimension in any anthropological inquiry. If biological anthropologists are interested in the identification of the genetic features distinguishing Pygmies from their neighbors, if they wish to reconstruct the population dynamics and the demographic processes accounting for the substantial genetic diversity between and within these two groups of populations, then the investigation of the cultural mechanisms accounting for the conservation of a significant cultural Pygmy specificity (as opposed to non-Pygmies with whom they are socially and economically largely tied) has a topical relevance. Bahuchet agrees that a better knowledge of the migratory flows of African Pygmies throughout Central Africa is needed since the early contacts with non-Pygmies would certainly provide more reliable micro-evolutionary insights on the genetic side. This is why addressing the historical relationship between languages spoken by Pygmies and non-Pygmies will provide a hypothesis that may be tested by a molecular approach.

Second, Bahuchet notes that even disciplines that are commonly perceived as distant from biological anthropology are worth exploring and merging with other lines of evidence. For instance, studying musical practices (e.g., Fürniss and Bahuchet 1995) across populations may provide important insights about past and present cultural contacts between populations. Such “alternative” approaches are essential for understanding the early contacts between Pygmies and non-Pygmies in the absence, we stress it once more, of archeological and demographic data for these populations.

Epilogue. By combining past and current knowledge regarding both evolutionary and cultural aspects, we have shown how the category “Pygmy” can be misleading when used to describe culturally, morphologically, and biologically heterogeneous populations. However, when appropriately used, one can successfully consider this same term to address numerous questions of fundamental importance for the understanding of human evolution in Central Africa. Caution invoked here is not a secondary aspect: It is imperative that biological anthropologists precisely define what they mean by the word “Pygmy.” As Serge Bahuchet convincingly explains, we should never forget that while Pygmies do not exist as a single homogeneous cultural group: populations such as the Aka,

Asua, Baka, Bongo, Efe, Kola, Koya, Sua, Twa, etc., exist as communities of people having rich, complex, and diverse cultures who are in contact with several neighboring ethnic groups with equally rich, complex, and diverse cultures. If such a proviso does apply to all human populations, we note that “biological anthropologists” often tend to use misleading terminologies, as some past investigations of Central African Pygmy and non-Pygmy populations have shown. Let us think about how many messages and inspirations these groups could give to us if we deconstructed more systematically the Western prism by which we perceive them, and if we tried to embrace and correct more openly our lack of understanding of the complexity of the Central African pluri-ethnic context.

Acknowledgments The authors would like to sincerely thank Dr. Franz Manni and Dr. Pierre Darlu for their many words of advice and numerous comments concerning this editorial. Paul Verdu wants to thank Evelyne Heyer and colleagues at the CNRS and the MNHN of Paris (UMR 7206 “Ecoanthropology and Ethnobiology”) for giving him the opportunity to work on Central African Pygmy populations between 2005 and 2008, a work which was supported by the ACI Prosodie. The current work is supported in part by the US National Institutes of Health grant RO1 GM081441.

Giovanni Destro Bisol is grateful to Gabriella Spedini for giving him the opportunity to study Pygmy populations from the Central African Republic and Cameroon. He was supported by grants from the University of Rome “La Sapienza,” the Ministero dell’Istruzione, dell’Università e della Ricerca, and the Istituto Italiano di Antropologia.

Received 14 September 2011; revision accepted for publication 13 October 2011.

Literature Cited

- Bahuchet, S. 1992. Spatial mobility and access to the resources among the African Pygmies. In *Mobility and Territoriality Social and Spatial Boundaries among Foragers, Fishers, Pastoralists and Peripatetics*, Casimir and Rao, eds. NY/Oxford: Berg, 205–257.
- Bahuchet, S. 1993. L’Invention des Pygmées. *Cahiers d’Etudes Africaines* 129 (Tome 33 n°1):153–181.
- Bahuchet, S. 2011. Changing Language, Remaining Pygmy. *Hum. Biol.* 84(1):11–43.
- Bahuchet S., and H. Guillaume. 1982. Aka-farmer relations in the northwest Congo basin. In *Politics and History in Band Societies*, E. Leacock and R. B. Lee, eds. Cambridge/Paris: Cambridge University Press/M.S.H., 189–211.
- Bahuchet, S., C. M. Hladik, A. Hladik, and E. Dounias. 1989. Les stratégies agricoles complémentaires de la chasse et de la pêche. In *Se Nourrir En Forêt Équatoriale: Anthropologie Alimentaire Des Populations Des Régions Forestières Humides d’Afrique*, C. M. Hladik, S. Bahuchet, and I. de Garine, eds. Paris: Unesco-MAB. 31–34.
- Bailey, R. C., G. Head, M. Tenike et al. 1989. Hunting and gathering in tropical rainforest: Is it possible? *Am. Anthropol.* 91:59–82.
- Batini, C., V. Coia, C. Battaglia et al. 2007. Phylogeography of the human mitochondrial L1c haplogroup: Genetic signatures of the prehistory of Central Africa. *Mol. Phylogenet. Evol.* 43(2):635–644.

- Batini, C., J. Lopes, D. M. Behar et al. 2011. Insights into the demographic history of African Pygmies from complete mitochondrial genomes. *Mol. Biol. Evol.* 28(2):1099–1110.
- Beaumont, M. A., W. Zhang, and D. J. Balding. 2002. Approximate Bayesian computation in population genetics. *Genetics* 162(4):2025–2035.
- Becker, N. S., P. Verdu, A. Froment et al. 2011. Indirect evidence for the genetic determination of short stature in African Pygmies. *Am. J. Phys. Anthropol.* 145(3):390–401.
- Becker, N. S. A., P. Verdu, B. Hewlett, and S. Pavard. 2010. Can life history trade-offs explain the evolution of short stature in human pygmies? A response to Migliano and colleagues. *Hum. Biol.* 82(1):17–27.
- Campbell, M. C., and S. A. Tishkoff. 2010. The evolution of human genetic and phenotypic variation in Africa. *Curr. Biol.* 20(4):R166–173.
- Cavalli-Sforza, L. L., ed. 1986. *African Pygmies*. London: Academic Press.
- Cornelissen, E. 2002. Human responses to changing environments in Central Africa between 40,000 and 12,000 B.P. *J. World Prehistory* 16(3):197–235.
- Cornuet, J. M., F. Santos, M. A. Beaumont et al. 2008. Inferring population history with DIY ABC: A user-friendly approach to Approximate Bayesian Computation. *Bioinformatics* 24(23):2713–2719.
- Destro-Bisol, G., V. Coia, I. Boschi et al. 2004a. The analysis of variation of mtDNA hypervariable region 1 suggests that Eastern and Western Pygmies diverged before the Bantu expansion. *Am. Nat.* 163(2):212–226.
- Destro-Bisol, G., F. Donati, V. Coia et al. 2004b. Variation of female and male lineages in sub-Saharan populations: The importance of sociocultural factors. *Mol. Biol. Evol.* 21(9):1673–1682.
- Diamond, J. M. 1991. Anthropology. Why are pygmies small? *Nature* 354(6349):111–112.
- Froment, A. 1993. Adaptation biologique et variation dans l'espèce humaine: Le cas des Pygmées d'Afrique. *Bull. Mém. Soc. Anthropol. Paris.* 5:417–448.
- Fürniss, S., and S. Bahuchet. 1995. Existe-t-il des instruments de musique pygmées? In *Ndroje Balendro, Musiques, Terrains Et Disciplines (Textes Offerts À Simha Arom)*, V. Dehoux, S. Fürniss, S. Le Bomin, E. Olivier, H. R. and F. Voisin, eds. Paris: Peters-Selaf. 87–109.
- Greenberg, J. H. 1966. *The Languages of Africa*. The Hague: Indiana University/Mouton & Co.
- Guthrie, M. 1967. *Comparative Bantu: An Introduction to the Comparative Linguistics and Prehistory of the Bantu Languages*. Farnborough: Gregg. 4 v. p.
- Hewlett, B. 1996. Cultural diversity among African Pygmies. In *Cultural Diversity among Twentieth-Century Foragers*. In *An African Perspective*, S. Kent, ed. Cambridge: Cambridge University Press, 215–244.
- Hiernaux, J. 1974. *The People of Africa*. London: Weidenfeld and Nicolson.
- Joiris, D. V. 2003. The framework of Central African hunter-gatherers and neighbouring societies. *African Study Monographs*, Suppl. 28:57–79.
- Kazadi, M. 1981. Méprisés et admirés: L'ambivalence des relations entre les Bacwa (Pygmées) et les Bahemba (Bantu). *Africa.* 51(4):837–847.
- Migliano, A. B., L. Vinicius, and M. M. Lahr. 2007. Life history trade-offs explain the evolution of human pygmies. *Proc. Natl. Acad. Sci. USA* 104(51):20216–20219.
- Patin, E., G. Laval, L. B. Barreiro et al. 2009. Inferring the demographic history of African farmers and pygmy hunter-gatherers using a multilocus resequencing data set. *PLoS Genet.* 5(4):e1000448.
- Perry, G. H., and N. J. Dominy. 2009. Evolution of the human pygmy phenotype. *Trends Ecol. Evol.* 24(4):218–225.
- Quintana-Murci, L., H. Quach, C. Harmant et al. 2008. Maternal traces of deep common ancestry and asymmetric gene flow between Pygmy hunter-gatherers and Bantu-speaking farmers. *Proc. Natl. Acad. Sci. USA* 105(5):1596–1601.
- Tishkoff, S. A., F. A. Reed, F. R. Friedlaender et al. 2009. The genetic structure and history of Africans and African Americans. *Science* 324(5930):1035–1044.

- Turnbull, C. M. 1965. *Wayward Servants, the Two Worlds of the African Pygmies*. New York: The Natural History Press.
- Verdu, P., F. Austerlitz, A. Estoup et al. 2009. Origins and genetic diversity of pygmy hunter-gatherers from Western Central Africa. *Curr. Biol.* 19(4):312–318.
- Weir, B. S., and C. C. Cockerham. 1984. Estimating F-statistics for the analysis of population-structure. *Evolution* 38(6):1358–1370.